

Drug Alcohol Depend, July 1, 2005; 79(1): 1-10.

Buprenorphine versus methadone in the treatment of pregnant opioid-dependent patients: effects on the neonatal abstinence syndrome.

HE Jones, RE Johnson, DR Jasinski, KE O'Grady, CA Chisholm, RE Choo, M Crocetti, R Dudas, C Harrow, MA Huestis, LM Jansson, M Lantz, BM Lester, and L Milio

The Department of Psychiatry and Behavioral Sciences, Johns Hopkins Bayview Medical Campus, Johns Hopkins University School of Medicine, D-3-East, 4940 Eastern Avenue, Baltimore, MD 21224, USA. hejones@jhmi.edu

MEDLINE ABSTRACT

This study was designed to compare the neonatal abstinence syndrome (NAS) in neonates of methadone and **buprenorphine** maintained pregnant opioid-dependent women and to provide preliminary safety and efficacy data for a larger multi-center trial. This randomized, double-blind, double-dummy, flexible dosing, parallel-group controlled trial was conducted in a comprehensive drug-treatment facility that included residential and ambulatory care. Participants were opioid-dependent pregnant women and their neonates. Treatment involved daily administration of either sublingual **buprenorphine** or oral methadone using flexible dosing of 4-24 mg or 20-100 mg, respectively. Primary a priori outcome measures were: (1) number of neonates treated for NAS; (2) amount of opioid agonist medication used to treat NAS; (3) length of neonatal hospitalization; and (4) peak NAS score. Two of 10 (20%) **buprenorphine**-exposed and 5 of 11 (45.5%) methadone-exposed neonates were treated for NAS ($p=.23$). Total amount of opioid-agonist medication administered to treat NAS in methadone-exposed neonates was three times greater than for **buprenorphine**-exposed neonates (93.1 versus 23.6; $p=.13$). Length of hospitalization was shorter for **buprenorphine**-exposed than for methadone-exposed neonates ($p=.021$). Peak NAS total scores did not significantly differ between groups ($p=.25$). Results suggest that **buprenorphine** is not inferior to methadone on outcome measures assessing NAS and maternal and neonatal safety when administered starting in the second trimester of pregnancy.